

Global Forum on Biological Control and Training Workshop on Biological Control

Nairobi, Kenya 26-30 June 2023

Studies on non-target effects of biopesticides

Evanson Omuse (MSc)

Co-organized by



Food and Agriculture
Organization of the
United Nations



icipe

Supported by



EUROPEAN UNION



USAID
FROM THE AMERICAN PEOPLE



Implemented by

giz Deutsche Gesellschaft
für Internationale
Zusammenarbeit (GIZ) GmbH

Introduction

❖ Agriculture partly depends on the beneficial services offered by nature:



- Over 75% of flowering crops depend on **insect pollinators**.
- Some pollinators (bees) provide hive products such as honey, wax, cerumen, bee bread, royal jelly, bee venom, and propolis.



- **Natural enemies** (parasitoids and predators) reduce pest populations.



- **Soil invertebrates** improve fertility.


Chemical insecticides constitute the key drivers to the **alarming global decline** of pollinators.



Food and Agriculture
Organization of the
United Nations



icipe



Dusts

Tree pollen

Foliar pesticide residue

Nectar

Pollen

Guttation water

Pesticide overspray

Propolis

honey,
wax,
cerumen,
bee bread,
royal jelly,
bee venom,
propolis

Systemic pesticides

Water



Do maize plants require insect pollination?

- Bees foraging on maize, other flowering crops (legumes, sunflowers) and wildflowers (weeds) on, and around maize field may be exposed to several chemical insecticides sprayed on fields (Fernandez-Cornejo et al. 2014; Long and Krupke 2016).
- To protect pollinators, biopesticides based on entomopathogenic fungi (EPF) can be used as alternatives to chemical insecticides.



Biopesticides developed at icipe



Ecotoxicological test of biopesticides



- Ecotoxicological dossiers are available for registered biopesticides.
- Ecotoxicological dossiers are based on Organisation for Economic Co-operation and Development (OECD, 1998) guidelines to test suboptimal doses ($<10^7$ conidia/mL) through oral exposure in short bioassays (<96 hr) with honeybees.
- Stingless bees have never been used as test insects.
- Long (10-days) bioassays with honeybees and stingless bees can result in variable toxicity levels depending on EPF isolates.
- International Organization of Biological Control (IOBC) classification:
 - ❖ Class 1: harmless ($<25\%$),
 - ❖ Class 2: slightly harmful ($25\%-50\%$),
 - ❖ Class 3: moderately harmful ($51\%-75\%$),
 - ❖ Class 4: harmful ($>75\%$)



Food and Agriculture
Organization of the
United Nations



Selected biopesticides screened for non-target effect on pollinators (honeybees and stingless bees)



Biopesticide isolate name	Trade Name	Target pests
<i>Metarhizium anisopliae</i> ICIPE 7	Detain	Fall armyworm
<i>Metarhizium anisopliae</i> ICIPE 78	Achieve, Mazao achieve	Fall armyworm, Spider mites,
<i>Metarhizium anisopliae</i> ICIPE 20	-	Tuta absoluta, Fall armyworm, leafminers
<i>Metarhizium anisopliae</i> ICIPE 62	Mazao supreme	Aphids
<i>Metarhizium anisopliae</i> ICIPE 69	Campaign	Thrips, whiteflies, leafminers
<i>Beauveria bassiana</i> ICIPE 284	-	Fall armyworm


Akutse et al., 2020



Food and Agriculture
Organization of the
United Nations



Objectives of the study



Effect biopesticides on
survival of bees in lab



Modelling performance of
the biopesticides in bee
colonies simulated
temperatures.



Effect of biopesticides on
bee pollination behaviour
and crop productivity in
semi-field trials



Effect of biopesticides on survival of bees

Honeybees
Apis mellifera



Stingless bees
Meliponula ferruginea



Spray tower



Fungal isolates

- Applying 10^8 conidia/mL or water on filter paper.
- Exposure of 25-30 bees/cage in four replicates.

Incubators

Honeybees



Stingless bees



10-days survival bioassays

Results

Kaplan-Meier survival curves of fungus-exposed bees



Honeybees

Stingless bees



1st experiment

2nd experiment



The tested isolates were moderately and **nontoxic** to bees according to the **International Organization of Biological Control (IOBC)** classification



Food and Agriculture
Organization of the
United Nations



Journal of Economic Entomology, 115(1), 2022, 46–55

<https://doi.org/10.1093/jee/toab211>

Advance Access Publication Date: 22 November 2021

Research



ENTOMOLOGICAL
SOCIETY OF AMERICA
SHARING INSECT SCIENCE GLOBALLY



Biological and Microbial Control

Susceptibility of the Western Honey Bee *Apis mellifera* and the African Stingless Bee *Meliponula ferruginea* (Hymenoptera: Apidae) to the Entomopathogenic Fungi *Metarhizium anisopliae* and *Beauveria bassiana*

Evanson R. Omuse,^{1,2,●} Saliou Niassy,^{1,●} John M. Wagacha,² George O. Ong'amo,² H. Michael G. Lattorff,^{1,●} Nkoba Kiatoko,¹ Samira A. Mohamed,¹ Sevgan Subramanian,^{1,●} Komivi S. Akutse,^{1,●} and Thomas Dubois^{1,3}

¹International Centre of Insect Physiology and Ecology (*icipe*), 30772-00100, Nairobi, Kenya, ²School of Biological Sciences, University of Nairobi, 30197-00100, Nairobi, Kenya, and ³Corresponding author, e-mail: tdubois@icipe.org

Subject Editor: Reed Johnson

Received 19 April 2021; Editorial decision 8 October 2021



Food and Agriculture
Organization of the
United Nations



Modelling the performance of the biopesticide isolates under bee colonies' temperatures




Conidial germination




Fungal suspension
(10^6 conidia/mL)

Mycelial growth




Mycelial mat

Incubation at
12, 16, 20, 24,
28, 32, 36°C



Observation of
conidial germ tubes



Measurement of
radial growth

Model selection

Eight models from
literature

Model validation

Output

Model
comparison

- Goodness-of-fit
- Adjusted R-squared

**Thermal requirements
for isolates:**
Tmin (°C), Topt (°C),
Tmax (°C), Pmax
(%germination, mm/day)



Food and Agriculture
Organization of the
United Nations



Temperature dependent models tested



Model tested

1. Briere 1
2. Briere 2
3. Ratkowsky 2
4. Ratkowsky 3
5. Lactin 1
6. Van Der Heide
7. CTMI (Cardinal temperature model with Inflection)
8. Generalized beta function




Food and Agriculture
Organization of the
United Nations




Results

Conidial germination curves by temperature-dependent models



Conidial germination **started** from 10–14°C and reached **a peak** at 25–30°C, followed by a precipitous decline reaching **upper thresholds** at 35–38°C.

Growth curves as predicted by 8 temperature-dependent models



Isolate	Tmin (°C)	Topt (°C)	Tmax (°C)	Pmax (mm/day)
ICPIPE 7	7.4	27.6	36.8	1.89
ICPIPE 20	13.5	26.8	37.0	2.28
ICPIPE 62	3.7	28.4	37.2	1.85
ICPIPE 69	8.7	27.1	36.6	1.89
ICPIPE 78	7.3	26.5	36.7	1.54
ICPIPE 284	9.5	25.9	35.4	1.36





Biocontrol Science and Technology



ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/cbst20>

Suitable models to describe the effect of temperature on conidial germination and mycelial growth of *Metarhizium anisopliae* and *Beauveria bassiana*

Evanson R. Omuse, Saliou Niassy, John M. Wagacha, George O. Ong'amo, Abdelmutalab G. A. Azrag & Thomas Dubois

To cite this article: Evanson R. Omuse, Saliou Niassy, John M. Wagacha, George O. Ong'amo, Abdelmutalab G. A. Azrag & Thomas Dubois (2021): Suitable models to describe the effect of temperature on conidial germination and mycelial growth of *Metarhizium anisopliae* and *Beauveria bassiana*, Biocontrol Science and Technology, DOI: [10.1080/09583157.2021.1993133](https://doi.org/10.1080/09583157.2021.1993133)


To link to this article: <https://doi.org/10.1080/09583157.2021.1993133>




Food and Agriculture
Organization of the
United Nations



Effect of a biopesticide on survival, pollination behaviour and success of the stingless bee




Cultivation of cucumber
in screenhouses




Experiment was repeated three times in Apr 2020 – Jun 2020, Sep 2020 – Nov 2020, and Dec 2020 – Feb 2021

Results

Foraging
activity




Survival of forager stingless bees



Fruit set and
maturation (%)

Days	Fruit set (%)			Mature fruits (%)	
	Control	Biopesticide		Control	Biopesticide
-9 – -7	88 ± 2	89 ± 3		80 ± 4	80 ± 4
-6 – -4	94 ± 1	91 ± 2		88 ± 2	87 ± 2
-3 – -1	94 ± 1	89 ± 5		90 ± 2	87 ± 2
0 – 2	94 ± 2	96 ± 3		84 ± 5	88 ± 5
3 – 5	94 ± 1	92 ± 2		89 ± 2	88 ± 2
6 – 8	88 ± 1	86 ± 6		81 ± 2	76 ± 2



Fruit weight (g)

Days	Apr 2020 – Jun 2020		Sep 2020 – Nov 2020		Dec 2020 – Feb 2021	
	Control	Biopesticide	Control	Biopesticide	Control	Biopesticide
-9 – -7	296 ± 15	257 ± 22	354 ± 13	358 ± 27	338 ± 15	371 ± 29
-6 – -4	289 ± 22	272 ± 22	366 ± 15	376 ± 21	363 ± 18	366 ± 21
-3 – -1	246 ± 22	234 ± 28	362 ± 36	359 ± 14	379 ± 42	372 ± 22
0 – 2	304 ± 21	242 ± 26	379 ± 19	353 ± 15	376 ± 24	380 ± 27
3 – 5	248 ± 20	253 ± 37	385 ± 83	336 ± 24	372 ± 22	378 ± 20
6 – 8	234 ± 24	196 ± 16	346 ± 31	311 ± 20	364 ± 20	376 ± 27



Food and Agriculture
Organization of the
United Nations



icipe

Apidologie (2022) 53:28







© The Author(s), 2022

<https://doi.org/10.1007/s13592-022-00938-1>

Original article



A fungal-based pesticide does not harm pollination service provided by the African stingless bee *Meliponula ferruginea* on cucumber (*Cucumis sativus*)

Evanson R. OMUSE^{1,2} , Saliou NIASSY¹ , Nkoba KIATOKO¹ , H. Michael G. LATTORFF¹ ,
John M. WAGACHA² , Thomas DUBOIS¹ 

¹ International Centre of Insect Physiology and Ecology (*icipe*), 30772–00100 Nairobi, Kenya

² School of Biological Sciences, University of Nairobi, 30197–00100 Nairobi, Kenya

Received 15 June 2021 – Revised 13 December 2021 – Accepted 18 December 2021




Food and Agriculture
Organization of the
United Nations



Conclusions

- Mortality due to biopesticides did not exceed 17.4% for honey bees or 11.0% for stingless bees.
- Biopesticides can germinate and grow optimally in beehive conditions.
- Biopesticide did not hamper bee pollination behaviour and success.
- The tested biopesticides can be safely used as to control insect pests in bee-resourced crop systems.
- The interactions of bee pollinators and biopesticides can also be limited by:
 1. careful timing of biopesticide application to avoid peak foraging periods.
 2. improving 'lure and infect' application techniques.



Acknowledgements



- Dr Thomas Dubois, Dr. Saliou Niassy, Dr. Subramanian Sevgan, Dr. Samira Mohamed.
- Technical team at Arthropod Pathology Unit – *icipe*.
- Funding sources.



Food and Agriculture
Organization of the
United Nations





Thank you

Co-organized by

Supported by

