



Global Forum on Biological Control and Training Workshop on Biological Control

Nairobi, Kenya 26-30 June 2023

Basic and Concept on microbial control – potential for FAW management

Komivi S. Akutse - kakutse@icipe.org

Co-organized by



Food and Agriculture
Organization of the
United Nations



icipe

Supported by



USAID
FROM THE AMERICAN PEOPLE




Implemented by

giz
Deutsche Gesellschaft
für Internationale
Zusammenarbeit (GIZ) GmbH


Estimated world biopesticide sales & alternative to pesticides

- Global chemical market is projected to grow from **\$70.24 bn** in 2019 to **\$101.64 bn** in 2026 ~ **5.42%**
- Biopesticides **\$4.4 billion** in 2019; **\$10.6 billion** by 2027; will match chemicals by 2050




\$404.5 M - in 2010

Predicted growth of Biopesticides sector



Source: Lux Research, Inc. www.luxresearchinc.com

PPPs to move new developed biopesticides in Africa



Arthropod Pathology Unit



- ❖ Research and innovations on biopesticides
- ❖ Bioprospecting for entomopathogens
- ❖ Microbes and pests' identification and characterization
- ❖ Maintains a long-term repository of entomopathogens
- ❖ Capacity-building in insect pathology research
- ❖ Commercialization in partnership with private sector



Food and Agriculture
Organization of the
United Nations



icipe

Global food security & why pests control?



A recent study showed that, insect invasions cost at least \$162.7 billion to the global economy & insects reduced the global agricultural production by 10- 16%

- ❖ While associated health costs exceed \$6.9 billion per year
- ❖ Invasive species are costing Africa \$3.66 trillion per year — *Nature* 592, 571-576 (2021)

Strategies used to overcome pest losses

- ✓ Increased & abusive use of pesticides/ insecticides (Costly, resistance development, health issues, residues accumulation, non target effects, etc.)
- ✓ Ecological approaches (Resistant host plants, Quarantine, Barriers, Trap crops, Crop diversity, etc.)
- ✓ GMO
- ✓ Cultural practices & Physical control and others...
- ✓ **Biological control** (Predators, Parasitoids, **Pathogens (biopesticides)** & Biorationals
- More sustainable, environmentally friendly, cost effective, etc.)



Food and Agriculture
Organization of the
United Nations



Invasion of FAW - Over dependence on pesticides



Business Guide Africa
Turning your Vision into Money

Home Hotels Flights Rental Cars Cruises Tours

The EastAfrican NEWS BUSINESS OPED SCIENCE & HEALTH MAGAZINE RWANDA TODAY

Rwanda faces hunger over fall armyworm invasion

SATURDAY APRIL 22 2017

Are there sustainable alternative approaches to pesticides? Yes, Nature regulate itself – How?



CNN Regions » Armyworm invasion destroys crops in southern Africa



Zambia is using its air force against the pests.



Food and Agriculture Organization of the United Nations



Ecosystem & entomopathogens



Ecosystem = community of living organisms in conjunction with the non-living components of their environment & interacting as a system

Ecosystem services = Benefits people obtain from ecosystems. The four categories of ecosystem services – *Supporting, Provisioning, Regulating & Cultural*
https://en.wikipedia.org/wiki/Ecosystem_services


Entomopathogens can also play diverse roles in nature including living as **Epiphytes** or **Endophytes** – microorganisms that colonize internal plant tissues without having any symptomatic effects on the host plant



Food and Agriculture
Organization of the
United Nations



Entomopathogens = Insect Pathogens



Fungi - *Beauveria bassiana*, *Metarhizium anisopliae*, *Hypocrea lixii*, *Isaria sinclairii*, *Lecanicillium spp.*, etc.

Bacteria - *Bacillus thuringiensis*: *Bt. kurstaki*, *Bt. aizawai*, *Bt. israelensis*, *Bt. tenebrionis*, *Bt. Japonensis*; *Serratia spp.* & *Paenibacillus popilliae*

Nematodes - *Steinernema carpocapsae*, *S. riobravis*, *S. scapterisci* & *Heterorhabditis bacteriophora*

Viruses - Nuclearpolyhedrosis virus (Japanese Beetle Milky Disease)



Epizootics or epidemics occurred in insect populations & we use these same causal agents as a strategy to control our insect pests ===== **Biopesticides development**

Entomopathogenic fungi (EPF) are considered as important BCAs, traditionally applied in inundative approach


However, they can also play diverse roles in nature including living as **Endophytes & Biofertilizers**

Beauveria bassiana

Gibberella sp.

Above & below ground effects of EPF –

“Lifestyle & ability to occupy diverse niches”



✓ Below- and aboveground communities of fungal entomopathogens are different

✓ *Metarhizium anisopliae* is most common belowground, but absent aboveground

✓ *Beauveria bassiana* is most commonly infecting arthropods aboveground

✓ *Metarhizium relyi* – known to infect a lot FAW



Metarhizium anisopliae




Food and Agriculture
Organization of the
United Nations



icipe

Strategic framework for development and evaluation of biopesticides



How do we value their service?



- ✓ **Bioprospecting** – Dead, sick, mycosis insects, host plants, soil...
 - ✓ **Isolations & Substrates selection** – Detection and isolation / culture and purification
 - ✓ **Identification** – Morphological and molecular tools
 - ✓ **Bioassays** – reinfection for virulence and efficacy against insect pest
 - ✓ **Formulations** – Aqueous & oil formulations (e.g. *Beauveria bassiana* & *Metarizhium anisopliae*)
 - ✓ **Virulence & Pathogenicity tests** – Conduct bioassays on target pests to characterize the efficacy of the pathogen
 - ✓ **Mass production & Applications** – Inundative & augmentative releases, Autodissemination devices (Pheromone or Lures/ attractants combined with the pathogens)
- Proper formulation of a fungal agent, requires a thorough understanding of not only the **life cycle of the pathogen, but also the biological aspects of both the fungus & the target host**



Fungi isolation technique through surface sterilization & insect bait



- ✓ **Soil** (soil suspension or insect bait: *Galleria mellonella* & *Tenebrio molitor*)
- ✓ **Host plants** (Endophytic fungi)
- ✓ **Seeds** (Seed-borne endophytes)
- ✓ **Insects** (Dead, Infected or Mycosed insects)



Beauveria bassiana
emerging from *Vicia faba* seeds



G. mellonella larva infested
by *M. anispliae*





Endophytic fungal colonies
emerging from plated plant tissue



G. mellonella larva infested by
B. bassiana



FAW diversity and gut microbes in Kenya



Diversity of fall armyworm, *Spodoptera frugiperda* and their gut bacterial community in Kenya

Joseph Gichuhi¹, Subramanian Sevgan¹, Fathiya Khamis¹, Johnnie Van den Berg², Hannalene du Plessis², Sunday Ekesi¹ and Jeremy K. Herren^{1,3}




Metarhizium rileyi has been frequently observed in some of the samples.

Epizootics of the *Metarhizium rileyi* are also observed regularly in Nairobi

Biopesticides development pathway – Innovation chain

Control and
gical Control



Basic science and concept design

Advanced knowledge

Applied research

Delivery of biopesticides – Innovation chain



Food and Agriculture
Organization of the
United Nations



icipe

Bioprospecting and maintenance of a repository of biopesticides



Entomopathogen group	No. of isolates	Genus
Entomopathogenic fungi	311	<i>Beauveria</i> , <i>Metarhizium</i> , <i>Verticillium</i> , <i>Isaria</i> , and others
Entomopathogenic bacteria	157	<i>Bacillus thuringiensis</i> , <i>Serratia marcescens</i> and others
Endophytes	10	<i>Hypocrea</i> , <i>Trichoderma</i> , <i>Clonostachys</i> , and <i>Bionecteria</i>
Entomopathogenic nematodes	2	<i>Heterorhabditis</i> and <i>Steinernema</i>
Microsporidian	3	<i>Nosema</i> , <i>Malamoeba</i> and <i>Johenrea locustae</i>
Baculoviruses	2	<i>Spodoptera littoralis</i> NPV and <i>S. exigua</i> NPV



Biopesticide research and product development



Plant pests

Whiteflies, *Liriomyza* leafminers, cereal stemborers, diamondback moth, African bollworm, red spider mites, aphids, thrips, fruit flies, pod-borers, pod suckers, storage beetles, false codling moth, fall armyworm and tomato leafminer

Animal pests and disease vectors

Ticks (*Rhipicephalus* sp., *Boophilus* sp., *Amblyoma* sp.) and tsetse flies (*Glossina* sp.)

Human disease vectors

Mosquitoes, other key vectors include tsetse flies, ticks, sand flies, fleas, black flies and triatomine bug

Parasites and microbes' screening for safety in insects for food & feed



Biopesticide products



**Metarhizium
62**

Aphids

**Metarhizium
78**

Mites

**Metarhizium
7**

Ticks

**Metarhizium
69**

Fruit flies
Thrips
Mealybugs



In partnership with *icipe*



Food and Agriculture Organization of the United Nations



Outreach for commercialized biopesticides



Registration status - Registered in 13 countries



Biological Control and
Biological Control





Food and Agriculture
Organization of the
United Nations



Scaling up of icipe' s commercialized biopesticides



Acreage of biopesticides use



Building private sector partnerships





Thank you

Co-organized by



Food and Agriculture
Organization of the
United Nations



icipe

Supported by



EUROPEAN UNION



USAID
FROM THE AMERICAN PEOPLE



german
cooperation
DEUTSCHE ZUSAMMENARBEIT

Implemented by

giz Deutsche Gesellschaft
für Internationale
Zusammenarbeit (GIZ) GmbH





Global Forum on Biological Control and Training Workshop on Biological Control

Nairobi, Kenya 26-30 June 2023

Prospects and opportunities for Biopesticides (EPF, bacteria, baculoviruses and endophytes) based management of FAW

Komivi S. Akutse - kakutse@icipe.org

Co-organized by



Food and Agriculture
Organization of the
United Nations



icipe

Supported by



EUROPEAN UNION



USAID
FROM THE AMERICAN PEOPLE



Implemented by

giz Deutsche Gesellschaft
für Internationale
Zusammenarbeit (GIZ) GmbH

Sustainable Fall armyworm IPM strategy for Africa



FAW-IPM Africa-specific, science-led, sustainable and integrated management of the fall armyworm



Foreign, Commonwealth & Development Office


FAW Biopesticides development & upscaling



Food and Agriculture Organization of the United Nations



Building on icipe' s biopesticide experience



Identification of effective biopesticides against the invasive FAW



Food and Agriculture
Organization of the
United Nations






Development of a biopesticide products for all stages for FAW



Insect viral diseases – eg. **Baculoviruses**



Insect bacterial diseases – eg. ***Bacillus thuringiensis***



Insect fungal diseases – eg. ***Metarhizium anisopliae***; ***Beauveria bassiana***





Food and Agriculture Organization of the United Nations



icipe

Efficacy of EPFs against FAW egg and newly emerged larvae





ORIGINAL CONTRIBUTION WILEY JOURNAL OF APPLIED ENTOMOLOGY

Ovicidal effects of entomopathogenic fungal isolates on the invasive Fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae)

Komivi Senyo Akutse | Jane Wanjiru Kimemia | Sunday Ekesi | Fathiya Mbarak Khamis | Odhiambo Levi Ombura | Sevgan Subramanian



Field efficacy trial with ICIPE 7 & ICIPE 78 – Embu, Kenya



Food and Agriculture Organization of the United Nations



Field Efficacy Trial in Kenya – incidence of damage caused by FAW




Met 7 & Met 78 have reduced the damage incidence equally to chemical by >60%

For severity of damage and yield

- ✓ No significant difference were observed for Cob width and length
- ✓ More yield were obtained in T2, T3 and T4 in Embu

****Registered – 2021 in Kenya; 2022 in Uganda and Tanzania**




T1- Control (Water only as application) ; **T2-** Met 7 (Oil formulation); **T3-** Met 78 (Oil formulation); **T4-** Chemical (Radiant); **T5-** Wood ash; **T6-** Mixit (Oil only as application)




Efficacy of EPF on FAW adults



Healthy



Fungus-infected



Combining insect pathogenic fungi and a pheromone trap for sustainable management of the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae)

Komivi S. Akutse^{a,*}, Fathiya M. Khamis^a, Felicitas C. Ambele^{a,b}, Jane W. Kimemia^a, Sunday Ekesi^a, Sevgan Subramanian^a





❖ *M. anisopliae* – ICIPe 315 & ICIPe 7 and *B. bassiana* – ICIPe 621 & ICIPe 676 caused 100% mortality




Food and Agriculture Organization of the United Nations



Effect of horizontal transmission of EPF inoculum as per FAW sex



*****None of the eggs hatched in the fungal treatments Vs. 100% hatchability in the control**



Combining insect pathogenic fungi and a pheromone trap for sustainable management of the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae)


Komivi S. Akutse^{a,*}, Fathiya M. Khamis^a, Felicitas C. Ambele^{a,b}, Jane W. Kimemia^a, Sunday Ekesi^a, Sevgan Subramanian^a



Food and Agriculture Organization of the United Nations



Development of biopesticides – Bt against FAW larvae



- ✓ Potent Bt isolates identified.
- ✓ Bt43, Bt39 and Bt50 holds promise with higher mortality and faster kill of FAW.
- ✓ Field efficacy studies are planned with the private sector partners.



Food and Agriculture Organization of the United Nations



icipe

Baculoviruses for FAW management in Africa

- ❖ Novel baculovirus “Fawligen” tested in Kenya
- ❖ Maize yield advantage of 1.5 t/ha over untreated control
- ❖ **Fawligen officially registered in Kenya (Feb 2021)**
- ❖ Community small-holder production being tested
- ❖ 95% of farmers willing to pay for biopesticide if available at an agro-dealer near to them, at a price comparable to a synthetic insecticide

Source: Ivan Rwomushana. Senior Scientist, Invasive Species Management. **CABI**





Food and Agriculture
Organization of the
United Nations



R4D support for enhancing product effectiveness, non-target effects & Capacity building




Non target effects on major FAW parasitoids – Direct effects



Cotesia icipe

Fungal isolates
ICIPE 7 caused the highest mortality rate at 81.40%



Fungal isolates	Mean mortality%	Larval mortality	Parasitism rates	Sex ratios (F:M)	Cotesia icipe LT ₅₀
<i>M. anisopliae</i> ICIPE 7	73.95±7.49a	55.25±6.74a	35.75±4.80b	2:1	2.3±0.03a
<i>M. anisopliae</i> ICIPE 78	33.63±6.63b	28.25±4.41b	62.00±5.02a	2:1	5.2±0.06c
<i>M. anisopliae</i> ICIPE 41	66.33±7.29a	53.75±3.91a	36.75±3.82b	2:1	2.8±0.04b
<i>B. bassiana</i> ICIPE 621	36.59±6.65b	31.00±5.29b	58.00±5.24a	2:1	5.0±0.06d

ICIPE 78 caused the highest (61.25%) FAW eggs mortality

High parasitism rates were obtained in ICIPE 78 and ICIPE 621 & ICIPE 7 and ICIPE 41 had lowest LT₅₀




Food and Agriculture Organization of the United Nations




Non target effects on major FAW parasitoids – Indirect effects




ICIPE 7 - *C. icipe*




ICIPE 41 - *C. icipe*




ICIPE 7 - *T. remus*




ICIPE 41 - *T. remus*




ICIPE 78 - *C. icipe*




ICIPE 621 - *C. icipe*



ICIPE 78 - *T. remus*



ICIPE 621 - *T. remus*



Cotesia icipe

Indirect application of the biopesticides at lower concentrations could be applied together with the parasitoids without affecting their performance

Telenomus remus



Food and Agriculture Organization of the United Nations



Efficacy of various oil formulations of *M. anisopliae* ICIPÉ 41 on FAW



Fungal isolate	Fungal formulation	Larvae cumulated mortality (%)	Lethal time 50% (LT ₅₀)
<i>Metarhizium anisopliae</i> ICIPÉ 41	Canola oil formulation	76.07 ± 6.43b	2.06 ± 0.15b
	Corn oil formulation	72.5 ± 5.58b	2.26 ± 0.36b
	Olive oil formulation	70.36 ± 6.66b	2.52 ± 0.43b
	Aqueous formulation	15.15 ± 2.86a	8.11 ± 1.53a



Food and Agriculture Organization of the United Nations



Effects of oil formulations of *M. anisopliae* ICIZE 41 on FAW parasitoids



Treatments	<i>Cotesia icipe</i>	<i>Telenomous remus</i>	<i>Trichogramma</i> spp.
Canola oil Formulation	82.5 ± 4.33a	21.3 ± 3.14b	26.3 ± 3.75b
Olive oil Formulation	52.5 ± 9.68b	15.0 ± 2.04b	23.8 ± 2.39b
Corn oil Formulation	12.5 ± 3.23c	2.5 ± 1.44a	8.8 ± 1.25a
Aqueous Formulation	52.5 ± 9.68b	11.3 ± 1.25b	23.8 ± 2.39b
Control	2.5 ± 1.44d	2.5 ± 1.44a	5.0 ± 2.04a

Mortality rates of the parasitoid species induced by indirect application of ICIZE 41 formulations


Parasitism rates obtained after indirect application of ICIZE 41 formulations

Parasitoids / Treatments	<i>Cotesia icipe</i>	<i>Telenomous remus</i>	<i>Trichogramma</i> spp.
Canola oil Formulation	31.5 ± 2.85a	30.2 ± 2.65a	33.5 ± 2.85a
Olive oil Formulation	44.1 ± 7.23ab	38.8 ± 8.43ab	40.0 ± 5.21ab
Corn oil Formulation	79.3 ± 2.94c	87.3 ± 2.50c	82.3 ± 2.34c
Aqueous Formulation	54.3 ± 4.12b	49.1 ± 4.42b	47.1 ± 7.23b
Control	84.3 ± 3.74c	87.3 ± 2.50c	85.2 ± 3.98c

Efficacy of ICIPE 41 on FAW and maize grain yield at Migori and Mbita



Treatments	% FAW larvae mortality	% Mycosis	Lethal time 50% ± SE
Migori			
Spinetoram-based pesticide/ Radiant	96.1 ± 2.5 a	0.0 ± 0.0 c	3.8 (3.80–3.90) b
Mazao Achieve®	73.0 ± 1.3 c	30.0 ± 0.7 b	5.1 (5.12–5.26) a
Corn oil formulation of <i>M. anisopliae</i> ICIPE 41	81.3 ± 2.6 b	70.0 ± 0.5 a	5.2 (5.22–5.32) a
Control	-	0.0 ± 0.0 c	-
Mbita			
Spinetoram-based pesticide/ Radiant	98.7 ± 1.3 a	0.0 ± 0.0 c	3.8 (3.80–3.88) b
Mazao Achive®	74.3 ± 2.5 c	25.0 ± 0.4 b	4.7 (4.70–4.82) a
Corn oil formulation of <i>M. anisopliae</i> ICIPE 41	83.6 ± 1.5 b	66.3 ± 0.7 a	4.6 (4.56–4.68) a
Control	-	0.0 ± 0.0 c	-



agronomy



Article

Performance of *Metarhizium anisopliae* Isolate ICIPE 41 in the Laboratory and Field in Comparison to Another Fungal Biopesticide and a Chemical Product to Sustainably Control the Invasive Fall Armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae)

Joseph Munywoki^{1,2}, Leonidah Kerubo Omosa³, Sevgan Subramanian¹, David Kupesa Mfuti¹, Ezekiel Mugendi Njeru², Vaderament-A. Nchiozem-Ngnitedem³ and Komivi Senyo Akutse^{1,*}




Food and Agriculture Organization of the United Nations



Cumulated larval mortality induced by different treatments and their lethal time 50% (LT₅₀) after applications



Host plants effects on efficacy of entomopathogenic fungal-based biopesticides in the management of Fall armyworm



On forage plants, *M. anisopliae* ICIPE 7 caused highest mortality of 82% to larvae fed on Napier grass, followed by 67% larvae fed on Panicum as compared to Maize and Brachiaria



Food and Agriculture Organization of the United Nations



Determining effectiveness of nature-based solutions to manage FAW


An on-station and on-farm validation trial conducted in Uganda

The treatments

- 1) M7,
- 2) M78,
- 3) Climate-smart PPT
- 4) M78 + PPT,
- 5) M7 + PPT,
- 6) Insecticide control,
- 7) PPT + insecticide,
- 8) Mono-cropped maize


Trial setup

- 1) Treatments replicated 3 times RCBD at the station and farmer as replicate on-station.
- 2) Treatment applied at 3, 6, and 9 WAP



The findings


- 1) The biopesticide treatments reduced the FAW infestation significantly compared with mono-cropped maize
- 2) The yield obtained from different treatments significantly differed from mono-cropped maize.
- 3) The combined effect of Biopesticides and PPT were more effective in enhancing the maize yield
- 4) If treatments were applied based on FAW abundance not with fixed dates, the impact of the biopesticide would have been better



Key points to consider in biopesticides or mycoinsecticides development



- ✓ Biology & Ecology of the pest & the biocontrol agent
- ✓ Interactions between the pest & other organisms – **Non target effects**
- ✓ Availability & Accessibility of the biocontrol agent
- ✓ Easy to formulate and apply & can be easily incorporated into IPM strategies
- ✓ Sustainable & environmentally friendly
- ✓ Low production cost & Cost-effective applications
- ✓ Pathogenic & virulent to the pest
- ✓ UV-tolerant isolates & Develop products that meet farmers' needs
- ✓ **Not detrimental to the non target agents / Natural enemies & Pollinators**
- ✓ Commercialization networks establishment
- ✓ Shelf life and storage conditions

Formulations, packaging, applications and factors that affect the viability or efficacy of EPF



- ✓ Temperature
- ✓ UV
- ✓ Humidity
- ✓ Some chemical lures are also not compatible with fungal viability and virulence
- Glycerin (0.1 %), nutrient agar (0.1% and molasses (0.5%) could be added to each formulation as protectants against UV light
- Ecological engineering could also be explored to overcome environmental factors
- Endophytes could overcome environmental abiotic challenges associated with EPF applications
- Genetic engineering provides useful strategies to either increase fungal virulence or enhance fungal resistance to different stress factors - (Virulence & Environmental stability)

Endophytes for PGP & FAW management




Endophytic colonization of maize seedlings through seed inoculation

Endophytic colonization of maize seedlings through foliar application





Endophytes colonization persistence






Endophytic persistence of *Hypocrea lixii* F3ST1, *Trichoderma harzianum* F2R41, *Trichoderma asperellum* M2RT4, *Trichoderma* spp. F2L41, and *Trichoderma atroviride* F2S21 5 weeks post-germination

Plant growth promotion parameters





- *Trichoderma atroviride* F2S21
- *Trichoderma asperellum* M2RT4
- *Trichoderma harzianum* F2R41
- *Trichoderma* spp. F2L41
- *Hypocrea lixii* F3ST1
- Control

Effect of endophytic colonization on maize seedling growth parameters. **(A)** Mean plant height. **(B)** Mean wet and dry shoot weight. **(C)** Mean leaf length. **(D)** Mean leaf width and **(E)** Mean leaf number.





Endophytes effects on reproduction traits and feeding/defoliation



- *Trichoderma atroviride* F2S21
- *Trichoderma asperellum* M2RT4
- *Trichoderma harzianum* F2R41
- *Trichoderma* spp. F2L41
- *Hypocrea lixii* F3ST1
- Control



Endophytes effects on pupation and adult emergence



Building capacity of production entomopathogenic fungi



Business incubation for small-scale farmers on biopesticide production
Arthropod Pathology Unit, *icipe*, 21-27 March 2020

Fall armyworm biopesticides
7 videos • 68 views • Last updated on Jan 26, 2021

Unlisted

ICIPE - International Centre of Insect Physiology and Ecology

SUBSCRIBED

- 1 Autoclaving (Arabic subtitles) - ICIPE - International Centre of Insect Physiology and Ecology - 4:10
- 2 Media preparation (Arabic subtitles) - ICIPE - International Centre of Insect Physiology and Ecology - 3:37
- 3 Pin isolation (Arabic subtitles) - ICIPE - International Centre of Insect Physiology and Ecology - 4:17
- 4 Plate preparation (Arabic subtitles) - ICIPE - International Centre of Insect Physiology and Ecology - 8:44
- 5 Pour plate technique (Arabic subtitles) - ICIPE - International Centre of Insect Physiology and Ecology - 5:23
- 6 Slide culture EPF for identification - ICIPE - International Centre of Insect Physiology and Ecology - 8:37
- 7 EPF mass production - ICIPE - International Centre of Insect Physiology and Ecology - 12:03

Video-tutorial on basic entomopathological procedures and fungus production



Training facility for small-scale production of entomopathogenic fungi



Expanding public-private-partnership for biopesticide scaling




- ❖ Interaction between - regulatory authorities from Tanzania, Ethiopia, Kenya, Uganda and East African Community; 4 Biopesticides companies; National partners and researchers

- ❖ Strengthen **external partnerships and collaborations** - from different part of the world



**Achieve®
&
Detain®**



Food and Agriculture
Organization of the
United Nations



Donor Acknowledgement

Direct Financial Support to *icipe* from:-





Thank you

Co-organized by



Food and Agriculture
Organization of the
United Nations



Supported by



USAID
FROM THE AMERICAN PEOPLE



Implemented by

giz Deutsche Gesellschaft
für Internationale
Zusammenarbeit (GIZ) GmbH

